


What is an opioid overdose?



Opioids can cause bad reactions that make your breathing slow or even stop. This can happen if your body can't handle the opioids that you take that day.

TO AVOID AN ACCIDENTAL OPIOID OVERDOSE:

- Do not to mix your opioids with alcohol, benzodiazepines (Xanax, Ativan, Klonopin, Valium), or medicines that make you sleepy.
- Be extra careful if you miss or change doses, feel ill, or start new medications.
- Get into drug treatment.
- Don't use alone.
- If you have a test strip for fentanyl, test your drugs first.

Now that you have naloxone...

Tell someone where it is and how to use it.

Common opioids include:

GENERIC	BRAND NAME
Hydrocodone	Vicodin, Lorcet, Lortab, Norco, Zohydro
Oxycodone	Percocet, Oxycontin, Roxicodone, Percodan
Morphine	MSContin, Kadian, Embeda, Avinza
Codeine	Tylenol with Codeine, TyCo, Tylenol #3
Fentanyl	Duragesic
Hydromorphone	Dilaudid
Oxymorphone	Opana
Meperidine	Demerol
Methadone	Dolophine, Methadone
Buprenorphine	Suboxone, Subutex, Zubsolv, Bunavail, Butrans

* Heroin is also an opioid.

For more information on local efforts to curb opioid misuse, call (530) 225-5468 (in emergency, call 911) or go to www.norxabuse.org



Shasta County
**Health & Human
Services Agency**

Opioid safety and how to use naloxone



A GUIDE FOR COMMUNITY MEMBERS



Shasta County
**Health & Human
Services Agency**

How to identify an opioid overdose:

Look for these common signs:

- The person won't wake up even if you shake them or say their name
- Breathing slows or even stops
- Lips and fingernails turn blue or gray
- Skin gets pale and clammy

In case of overdose:

- 1 Call 911 and give naloxone**
If no reaction in 3 minutes, give second naloxone dose
- 2 Do rescue breathing or chest compressions**
Follow 911 dispatcher instructions
- 3 After naloxone**
Stay with person for at least 3 hours or until help arrives

NARCAN[®] (naloxone HCl) **NASAL SPRAY**

QUICK START GUIDE Opioid Overdose Response Instructions

Use NARCAN Nasal Spray (naloxone hydrochloride) for known or suspected opioid overdose in adults and children.

Important: For use in the nose only.

Do not remove or test the NARCAN Nasal Spray until ready to use.

1 Identify Opioid Overdose and Check for Response

Ask person if he or she is okay and shout name.

Shake shoulders and firmly rub the middle of their chest.

Check for signs of opioid overdose:

- Will not wake up or respond to your voice or touch
 - Breathing is very slow, irregular, or has stopped
 - Center part of their eye is very small, sometimes called "pinpoint pupils"
- Lay the person on their back to receive a dose of NARCAN Nasal Spray.



2 Give NARCAN Nasal Spray

Remove NARCAN Nasal Spray from the box.

Peel back the tab with the circle to open the NARCAN Nasal Spray.

Hold the NARCAN nasal spray with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.

Gently insert the tip of the nozzle into either nostril.

- Tilt the person's head back and provide support under the neck with your hand. Gently insert the tip of the nozzle into one nostril, until your fingers on either side of the nozzle are against the bottom of the person's nose.

Press the plunger firmly to give the dose of NARCAN Nasal Spray.

- Remove the NARCAN Nasal Spray from the nostril after giving the dose.

Get emergency medical help right away.

Move the person on their side (recovery position) after giving NARCAN Nasal Spray.

Watch the person closely.

If the person does not respond by waking up, to voice or touch, or breathing normally another dose may be given. NARCAN Nasal Spray may be dosed every 2 to 3 minutes, if available.

Repeat Step 2 using a new NARCAN Nasal Spray to give another dose in the other nostril. If additional NARCAN Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.

